

Recombinant Rat Interleukin-18 (rr IL-18)

Synonyms: Interferon-gamma-inducing factor, IGIF, IL-1g, IL-18, IL1F4, MGC12320, IFN-gamma-inducing factor, Interleukin-1 gamma, IL-1 gamma, lboctadekin, Il18.

Introduction: IL-18 is a proinflammatory cytokine. This cytokine can induce the IFN-gamma production of T cells. The combination of this cytokine and IL12 has been shown to inhibit IL4 dependent IgE and IgG1 production and enhance IgG2a production of B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine and thus negatively regulate its biological activity.

Description: Recombinant rat IL-18 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 158 amino acids and having a molecular mass of 18.26 kDa. The rat IL-18 purified by proprietary chromatographic techniques.

Source: *Escherichia Coli*.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation: The protein was lyophilized from a concentrated (1 mg/ml) solution in water containing 0.4x PBS pH=7.2.

Solubility: It is recommended to reconstitute the lyophilized rr IL-18 in sterile water not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability: Lyophilized rr IL-18 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution rr IL-18 should be stored for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity: Greater than 95.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be His-Phe-Gly-Arg-Leu.

Biological Activity: The ED50 as determined by the dose-dependant stimulation of IFN-gamma by murine lymph node cells (4×10^5 cells/well) co-stimulated with sub optimal dose of ConA (0.75µg/ml) is 12 ng/ml. The medium was RPMI containing 10% FCS and 50µM 2-ME.

Protein content: Protein quantitation was carried out by two independent methods:

1. UV spectroscopy at 280 nm using the absorbency value of 0.213 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics).
2. Analysis by RP-HPLC, using standard solution of IL-18 as a Reference Standard.

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

<i>small</i>	5 µg	Cat.N°	13340180
<i>medium</i>	25 µg	Cat.N°	13340184
<i>large</i>	100 µg	Cat.N°	13340186

ImmunoTools Excellent Quality - Advantageously priced