

anti-human CD16 PerCP-conjugated

PerCP - conjugated monoclonal antibody LNK16 to human CD16

Cat-No: **21279165**

500 µl

Clone: LNK16

Specificity: The antibody LNK16 reacts with CD16, a low affinity receptor for aggregated IgG (FcγRIII antigen). CD16 exists in two different isoforms: CD16a (FcγRIIIA; 50-65 kDa; expressed on NK-cells, monocytes and macrophages) and CD16b (FcγRIIIB; 48 kDa; mainly expressed on neutrophils).

HLDA V; WS Code M MA069

HLDA V; WS Code NK NK50

Immunogen: Normal human peripheral blood granulocytes

Isotype subclass: Mouse IgG1

Form: The purified antibody is conjugated with Peridin Chlorophyll under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing BSA and 15 mM sodium azide (pH 7.4)

Expiration date: The reagent is stable until the expiry date stated on the vial label

Storage conditions: Store at 4 °C. Avoid prolonged exposure to light

Application: Flow Cytometry

Background: CD16 (FcγRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcγRIII is expressed in two forms – FcγRIII-A and -B. FcγRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcεRI-γ subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcγRIII-A is associated, moreover, with FcεRI-β subunit. Besides IgG, FcεRI-A can be triggered also by oligomeric IgE. FcγRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

References: Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

Boyle JJ Human macrophages kill human mesangial cells by Fas-L-induced apoptosis when triggered by antibody via CD 16. Clin Exp Immunol 137:529-37 (2004)

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

ImmunoTools GmbH - Altenoyther Str. 10; 26169 Friesoythe; Germany
Tel +49-(0)4491-400997, Fax +49-(0)4491-400998, info@immunotools.com
www.immunotools.com